

From: Flatwoods Citrus

Vol. 15, No. 6

June 2012

Previous issues of the Flatwoods Citrus newsletter can be found at:
<http://citrusagents.ifas.ufl.edu/agents/zekri/index.htm>

FLOODING INJURY

Both surface and subsoil drainage is necessary to obtain adequate root systems for the trees.

Roots, like the rest of the tree, require oxygen for respiration and growth. Soils in Florida typically contain 20-21 % oxygen. When flooding occurs, the soil oxygen is replaced by water. This condition causes tremendous changes in the types of organisms present in the soil and in the soil chemistry.

Flooding injury would be expected if the root zone were saturated for 3 days or more during extended summer rains at relatively high soil temperatures (86-95° F). The rate of oxygen loss from the soil is much greater at high than at low temperatures. The potential for damage to roots is less obvious but equally serious when the water table is just below the surface. Flooding stress is much less when water is moving than when water is stagnant. The use of observation wells is a very reliable method for evaluating water-saturated zones in sites subject to chronic flooding injury.



Short-term estimates of flooding stress can be obtained by digging into the soil and smelling soil and root samples. Sour odors indicate an oxygen deficient environment. The presence of hydrogen sulfide (a disagreeable rotten egg odor) and sloughing roots indicate that feeder roots are dying. Under flooded conditions, root death is not exclusively associated with oxygen deficiency. Anaerobic bacteria (the kind that can grow only in the absence of oxygen) develop rapidly in flooded soils and contribute to the destruction of citrus roots. Toxic sulfides and nitrites formed by anaerobic sulfate- and nitrate-reducing bacteria are found in poorly drained groves. Sulfate-reducing bacteria require both energy and sulfates in order to change sulfates to sulfides. The best sources of energy have been found to be certain organic acids contained in citrus roots, grass roots, and buried pieces of

palmetto. Thus, citrus roots can contribute to their own destruction by being an energy source for these bacteria.

Symptoms of flooding injury may occur within a few days or weeks, but usually show up after the water table has dropped and the roots become stranded in dry soils. Leaf wilting, leaf drop, dieback, and chlorosis patterns may develop and tree death may occur. Trees subjected to chronic flooding damage are stunted with sparse canopies, dull colored, small leaves and produce low yields of small fruit. New flushes of growth will have small, pale leaves due to poor nitrogen uptake by restricted root systems. Usually, the entire grove is not affected, but most likely smaller more defined areas will exhibit the symptoms. Striking differences in tree condition can appear within short distances associated with only slight changes in rooting depths. Water damage may also be recognized by a marked absence of feeder roots and root bark, which is soft and easily sloughed.

With acute water damage, foliage wilts suddenly followed by heavy leaf drop. Trees may totally defoliate and actually die, but more frequently partial defoliation is followed by some recovery. However, such trees remain in a state of decline and are very susceptible to drought when the dry season arrives because of the shallow, restricted, root systems. Moreover, waterlogged soil conditions, besides debilitating the tree, are conducive to the proliferation of soil-borne fungi such as *Phytophthora* root and foot rot. These organisms cause extensive tree death especially in poorly drained soils.

Water damage may usually be distinguished from other types of decline by a study of the history of soil water conditions in the affected areas. Areas showing water damage are usually localized and do not increase in size progressively as do areas of spreading decline. Foot or root rot symptoms include a pronounced chlorosis of the leaf

veins caused by root damage and girdling of the trunk. Lesions also appear on the trunk usually near the soil level (foot rot) or roots die and slough-off (root rot). Flood damage does not produce lesions. Trees with blight or CTV are usually randomly distributed within the grove and diagnostic tests are available to distinguish them from water-damaged trees.

Citrus trees respond physiologically to flooding long before morphological symptoms or yield reductions appear. Photosynthesis and transpiration decrease within 24 hours of flooding and remain low as flooding persists. Water uptake is also reduced which eventually translates to decreased shoot growth and yields.

It is both difficult and costly to improve drainage in existing groves, so drainage problems should be eliminated when the grove area is prepared for planting by including a system of ditches, beds and/or tiling. Growers should not depend on the slight and often unpredictable differences in rootstock tolerance to waterlogging to enable trees to perform satisfactorily under such conditions. Trees, irrespective of scion and rootstock cultivars, should be planted under the best drainage conditions possible. Drainage ditches should be kept free of obstruction through a good maintenance program including chemical weed control. Tree recovery from temporary flooding is more likely to occur under good drainage structure maintenance conditions.

Do not disk a grove if trees were injured by flooding. Irrigation amounts should be reduced, but frequencies should be increased to adequately provide water to the depleted, shallow root systems. Soil and root conditions should be evaluated after the flooding has subsided. Potential for fungal invasion should be determined through soil sampling and propagule counts. If there is a *Phytophthora* problem, the use of certain fungicides can improve the situation.